

# The velocity of the photons

## Radius and the propagation speed of photons

### "Quantum foam"

(These article are protected by copyright, officially registered in IGAC under N.º 1195/2010)

#### Abstract

The time delays in the arrival time of photons of high energy gamma-ray bursts relative to the smaller energy photon, implies that the speed of light is not constant.

Several conclusions can be drawn from this observation.

Different velocities of photons leads, the different velocities for the light, so we can say that the velocity of light is not constant.

We also know that this is potential constant  $C^2$  which participates in the energy and that is constant for all forms of energy.

The light, or photons, by their nature twilight should travel at velocity slightly less than  $C$ , because if it were otherwise, they would only energy, and lose their corpuscular nature and as such no longer be photons.

Now, we know that the velocity of the most energetic photons moving at a lower velocity from  $C$ , so the potential energy of the photon interferes negatively with their movement.

This has caused discussion and controversy in the scientific community.

It is the search for an answer to why this article.

#### Introduction

Let's start with a syllogism.

Today we know that light, the photons of different energies propagate at different speeds.

We also know that the energy of matter depends on the speed that  $C$  is constant for all masses.

The speed of light, which is the photon, is not the same for all photons, so the speed of light is not

constant.

If the speed of light is not constant for all photons, then the speed of light is not C because it is constant.

## The velocity C

As we see below C is the speed of propagation of the radiation mass  $mC^2$  and C is the speed involved in the energy of mass.

The light, or photons, by their corpuscular nature should travel at speeds slightly less than C, because if it were otherwise, they would only energy, and lose their corpuscular nature.

Experience has proved the different speeds at moving photons, a phenomenon known generically as "quantum foam".

Speed C is therefore the maximum speed allowed in the universe, in any direction.

Therefore, we are faced with a maximum universal escape potential.

As  $\rho_u$ - density potential energy at our local.

$$C^2 = 2 G \rho_u$$

It is this potential that generates the energy of matter.

$$mC^2 = 2 G m \rho_u$$

## Photons

Photons we have the same principle.

On the surface the photon we find:

$\rho_u$  - The density of potential energy of the local where the photon moves.

$\rho_f$  - The energy density generated by photon.

The density of potential energy surface of the photon is given by:

$$\rho_o = \rho_u + \rho_f$$

$$mC^2 = 2 G m \rho_o$$

$$mC^2 = 2 G m (\rho_u + \rho_f)$$

Dividing both sides by  $\underline{m}$  why we are to make reference to the photon  $f$ .

$$C_f^2 = 2 G \rho_u + 2 G \rho_f$$

$$C_f^2 = C^2 + 2 G \rho_f$$

$$C_f^2 > C^2$$

This inequality can't happen, given the definition of mass energy  $mC^2$ , where in order not to violate the principle:

$$:K = \frac{C_f^2}{C^2}$$

$$K = \frac{2 G \rho_u + 2 G \rho_f}{2 G \rho_u}$$

$$K = \frac{\rho_u + \rho_f}{\rho_u}$$

K is the transformation of the G value of another benchmark for our reference from the quantities measured from our reference, provided in relativity RF.

$$\frac{G_f}{G_u} = \frac{\rho_u}{\rho_u + \rho_f} = \frac{t_f^2}{t_u^2}$$

Read on: [Remaking of the theory of relativity.Not bendof space.10/09](#)

$$C_f^2 = C^2 \frac{\rho_u + \rho_f}{\rho_u}$$

$$C^2 \frac{\rho_u + \rho_f}{\rho_u} = 2 G \rho_u + 2 G \rho_f$$

$$C^2 = 2 G \rho_u \frac{\rho_u}{\rho_u + \rho_f} + 2 G \rho_f \frac{\rho_u}{\rho_u + \rho_f}$$

$$C^2 = C^2 \frac{\rho_u}{\rho_u + \rho_f} + C^2 \frac{\rho_f}{\rho_u + \rho_f}$$

There are therefore two parts of a different nature in the creation of photon energy.

$m C^2 \frac{\rho_u}{\rho_u + \rho_f}$  - The energy density induced by the universal potential energy, causing the movement of

photons, ie their energy of motion or kinetic.

$m C^2 \frac{\rho_f}{\rho_u + \rho_f}$  - The intrinsic energy to photon energy density induced by the potential energy of the

photon itself.

The motion of photons is generated solely by the density of potential energy in the universal local he goes through and not by its own potential energy.

We can then represent the photon energy in the form:

The kinetic energy of the photon is given by ( $mV^2$ ).

$$mC^2 = mV^2 + 2Gm\rho_f \frac{\rho_u}{\rho_u + \rho_f}$$

$$C^2 = V^2 + C^2 \frac{\rho_u}{\rho_u + \rho_f} + C^2 \frac{\rho_f}{\rho_u + \rho_f}$$

$$C^2 = V^2 + C^2 \frac{\rho_f}{\rho_u + \rho_f}$$

**The speed of displacement of the photon:**

**The speed of displacement of the photon is given by.**

$$V^2 = C^2 - C^2 \frac{\rho_f}{\rho_u + \rho_f}$$

$$V = C \sqrt{1 - \frac{\rho_f}{\rho_u + \rho_f}}$$

$$V = C \sqrt{\frac{\rho_u}{\rho_u + \rho_f}}$$

**The variation of the speed of photons for C ,dV, is given by**

$$C^2 = V^2 + dV^2$$

$$dV^2 = C^2 - V^2$$

$$dV^2 = C^2 - C^2 \frac{\rho_u}{\rho_u + \rho_f}$$

$$dV^2 = C^2 \left( 1 - \frac{\rho_u}{\rho_u + \rho_f} \right)$$

$$dV^2 = C^2 \left( \frac{\rho_f}{\rho_u + \rho_f} \right)$$

$$dV = C \sqrt{\frac{\rho_f}{\rho_u + \rho_f}}$$

The value of the speed deviation of the photon for C is the escape potential of the photon itself.

Now, we know the speed of photons.

Increased energy density of photons ( $\frac{m}{r}$ ), lead to a slower velocity photons.

Actually C is the speed of propagation of radiation mass ( $\frac{m}{r}$ ) and not the speed of propagation of photons.

Photons of very low energy move very close to the speed C.

## Energy of particles

As we said before the energy of particles, such as the photon, is given by:

$\rho_m$  - Density of potential energy of matter.

$$mC^2 = 2 G m \rho_u \frac{\rho_u}{\rho_u + \rho_m} + 2 G m \rho_m \frac{\rho_u}{\rho_u + \rho_m}$$

$$mC^2 = mC^2 \frac{\rho_u}{\rho_u + \rho_m} + mC^2 \frac{\rho_m}{\rho_u + \rho_m}$$

That is the energy of matter is made up of two portions of different nature.

-  $mC^2 \frac{\rho_u}{\rho_u + \rho_m}$  The potential kinetic energy given by the density of potential energy universal.

-  $mC^2 \frac{\rho_m}{\rho_u + \rho_m}$  The intrinsic energy to matter itself

These are the readings on our referential.

Porto, 2/3/ 2010.

José Luís Pereira Rebelo Fernandes